

CHIRAL PERTUBATION THEORY WITH LATTICE REGULARIZATION.

I.A. Shushpanov

and

A.V. Smilga

*Institute for Theoretical and Experimental Physics, B. Cheremushkinskaya 25, Moscow
117218, Russia*

June 1998

Abstract

We study the structure and perform some explicit calculations for the power ultraviolet divergences and their renormalization in the Chiral Perturbation Theory with lattice regularization in 1 and 2 loops.

1 Introduction.

Chiral Perturbation Theory (ChPT) is nowadays a standard tool to study low-energy hadron dynamics [1]. The lagrangian of ChPT involves only lightest pseudoscalar degrees of freedom. The hadron excitations with higher mass appear only indirectly, via phenomenological constants of ChPT. As is well known, QCD enjoys an approximate $SU_L(N_f) \otimes SU_R(N_f)$ flavor chiral symmetry. This symmetry is exact in the limit of massless quarks. When quark masses are not zero, this symmetry is broken, but, while the masses are small, this breaking is under control and can be taken into account by a systematic expansion in mass. The lagrangian of ChPT is written in terms of the unitary matrix U which is usually parameterized as $U = \exp\{2i\phi^a t^a / F_\pi\}$ (ϕ^a are the pseudoscalar meson fields and F_π is the pion decay coupling constant) and involves all terms which are invariant under the chiral symmetry $SU_L(N_f) \otimes SU_R(N_f)$ and the non-invariant terms including quark mass matrix $\mathcal{M}$ and its higher powers. The theory is not renormalizable and involves an infinite number of counterterms which can be ordered by powers of derivatives (i.e. characteristic momenta) and quark mass. To the leading order

$$L^{(2)} = \frac{F^2}{4} \text{Tr}\{\partial_\mu U^\dagger \partial_\mu U\} + \Sigma \text{ReTr}\{\mathcal{M} U^\dagger\} , \quad (1)$$

where $\Sigma = | \langle \bar{q}q \rangle_{\text{vac}} |$ is the quark condensate. The lagrangian (1) involves a dimensional coupling $F \equiv F_\pi$ resulting in power ultraviolet divergences in the loops. The only way to get rid of them is to take into account counterterms with higher derivatives and higher powers of $\mathcal{M}$ (this is what non-renormalizability means). To perform the renormalization in practice, we have first to regularize the divergent integrals in ultraviolet. A naive momentum cut-off does not work here; it breaks chiral invariance explicitly. An almost exclusive way to handle the theory used so far in the literature is the dimensional regularization. It is practically quite convenient but has a drawback of being artificial and unphysical. In particular, it does not display power ultraviolet divergences (which are there in any physical scheme) whatsoever.

In this paper, we explore the lattice regularization scheme. As the only way to define what quantum theory actually means is to put it on the lattice (so that path integrals involve a finite number of variables), this regularization is the most natural and the most physical one from the "philosophical" viewpoint. In contrast to the dimensional regularization, it does involve power ultraviolet divergences explicitly which we consider as an advantage rather than a drawback. The price to be paid is the absence of Lorentz invariance in the regularized theory which makes the calculations somewhat more tricky than with dimensional regularization. But this technical complication is not really so serious.

Intermittently, the issue of lattice regularization and the related issue of the effects due to nontrivial measure in the path integral for the chiral theory was addressed in the literature [2], but we are not aware of any explicit calculation with this scheme. We will perform here some calculations with lattice regularization in 1 and 2 loops. For simplicity, we will assume $\mathcal{M}$ to be diagonal $\mathcal{M} = m\hat{1}$ and restrict ourselves to the case $N_f = 2$.

2 One-loop calculations.

Lattice version of the Euclidean action, corresponding to the chiral lagrangian (1) is following:

$$S = \frac{F^2 a^2}{2} \sum_{n,\mu} \text{Tr}\{1 - U_{n+e_\mu}^+ U_n\} - m \Sigma a^4 \sum_n \text{Tr}\{U_n\} , \quad (2)$$

where a is the lattice spacing and e_μ are the unit four-vectors connecting the adjacent nodes [writing the action (2), we used the fact that $\text{Tr}\{U\}$ is real when $U \in SU(2)$]. Many different parameterizations of the matrix U can be chosen. To begin with, we choose the Weinberg parametrization:

$$U = U_\mu \tau_\mu \quad [\tau_\mu = (1, i\tau^a); \quad U_\mu^2 = 1; \quad U^a = \pi^a/F] . \quad (3)$$

We may expand then lattice action (2) in the pion fields π^a . For 1-loop calculations, it suffices to restrict oneself by the terms $\sim \pi^2 \equiv \pi^a \pi^a$ and $\sim \pi^4$:

$$S = \sum_{n,\mu} \left[\frac{a^2}{2} (\pi_{n+e_\mu}^a - \pi_n^a)^2 + \frac{a^2}{4F^2} (\pi_n^4 - \pi_{n+e_\mu}^2 \pi_n^2) \right] + \frac{M_0^2 a^4}{2} \sum_n \left(\pi_n^2 + \frac{1}{4F^2} \pi_n^4 \right) , \quad (4)$$




Figure 1: One-loop graph for $\Pi(k)$.

where $M_0^2 = 2m\Sigma/F^2$ is the tree-level pion mass.

To the order $1/F^2$ there is only one relevant graph depicted in Fig. 1

To calculate it, we need to know the free scalar propagator on the lattice which can be taken e.g. from Ref.[3] :

$$D(x-y) = \int_{-\pi/a}^{\pi/a} \frac{d^4 q}{(2\pi)^4} \frac{e^{iq(x-y)}}{M_0^2 + \frac{2}{a^2} \sum_{\mu} [1 - \cos(q_{\mu}a)]} , \quad (5)$$

where $x_{\mu} = n_{\mu}a$, $y_{\mu} = n'_{\mu}a$ are the positions of lattice nodes and the momentum q_{μ} is restricted to lie in the interval $|q_{\mu}| \leq \pi/a$.

Calculating the polarization operator in Fig. 1, we obtain

$$\Pi^{\text{Fig.1}}(k) = \frac{2}{a^2 F^2} \left\{ \sum_{\mu} [D(0) - \cos(k_{\mu}a) D(ae_{\mu})] + \frac{5}{4} M_0^2 a^2 D(0) \right\} \quad (6)$$

where

$$D(0) = \int_{-\pi/a}^{\pi/a} \frac{d^4 q}{(2\pi)^4} \frac{1}{M_0^2 + \frac{2}{a^2} \sum_{\nu} (1 - \cos(q_{\nu}a))} , \quad (7)$$

$$D(ae_{\mu}) = \int_{-\pi/a}^{\pi/a} \frac{d^4 q}{(2\pi)^4} \frac{\cos(q_{\mu}a)}{M_0^2 + \frac{2}{a^2} \sum_{\nu} (1 - \cos(q_{\nu}a))} . \quad (8)$$

Here $D(ae_{\mu})$ is the propagator connecting adjacent lattice nodes. It is the same in all directions. When $a \ll M_0^{-1}$,

$$\begin{aligned} D(0) &\sim \frac{I}{a^2} , \\ D(ae_{\mu}) &\sim \frac{1}{a^2} \left(I - \frac{1}{8} \right) \end{aligned} \quad (9)$$

where

$$I = \frac{1}{2} \int_{-\pi}^{\pi} \frac{d^4 x}{(2\pi)^4} \frac{1}{\sum_i (1 - \cos x_i)} = \frac{1}{2} \int_0^{\infty} ds e^{-4s} I_0^4(s) \approx 0.154934 \quad (10)$$

[$I_0(s) = 1/2\pi \int_{-\pi}^{\pi} e^{s \cos x} dx$ is a Bessel function].

Eq.(6) is not, however, the final result for the polarization operator. Actually, it would be a disaster if it would. In the chiral limit $M_0 = 0$, $\Pi^{\text{Fig.1}}(k=0) = 1/(a^4 F^2) \neq 0$.

That would mean that the massless on the tree level pions acquire a mass (involving a quartic ultraviolet divergence) after renormalization. That would break chiral invariance [manifest in the lattice action (2)] which looks like a nonsense. The paradox is resolved by taking into account the fact that the path integral in chiral theory contains the integration over $SU(2)$ matrices U with the invariant Haar measure (otherwise the quantum theory would not be chirally invariant, indeed). We have:

$$\int [dU] = \int \Pi_{n,a} \frac{dU_n^a}{U_n^0} \sim \int \Pi_{n,a} \frac{dU_n^a}{1 - \frac{1}{2}U_n^2} \sim \int \Pi_{n,a} d\pi_n^a \exp \left\{ - \sum_n \ln \left(1 - \frac{1}{2}\pi_n^2/F^2 \right) \right\} . \quad (11)$$

Expanding logarithm, we obtain the extra term $-\frac{1}{2F^2} \sum_n \pi_n^2$ in the effective lattice action and, correspondingly, the following contribution in $\Pi(k)$

$$\Delta\Pi(k)[\text{measure}] = -\frac{1}{a^4 F^2} \quad (12)$$

[in the continuum limit, the contribution from the measure has the form $\sim F^{-2}\delta^{(4)}(0)$]. We see that the quartic divergent additive contributions to the pion mass coming from the 1-loop graph in Fig.1 and those from the measure cancel out completely.

The quadratic ultraviolet divergences in $\Pi(k)$ survive, however. Adding the contributions (6) and (12) and retaining the leading terms, we obtain

$$\Pi(k) = \frac{1}{a^2 F^2} \left[\frac{3IM_0^2}{2} + k^2 \left(I - \frac{1}{8} \right) \right] + o\left(\frac{1}{a^2}\right) . \quad (13)$$

We can now go over into the Minkowski space (with $k_E^2 \rightarrow -k_M^2$, $\Pi_E \rightarrow -\Pi_M$) and find the renormalized value of the mass as the solution to the dispersive equation $k_M^2 = M_0^2 + \Pi_M(k_M^2)$. We obtain

$$M_{phys}^2 = M_0^2 \left[1 + \frac{1}{2F^2 a^2} \left(I + \frac{1}{4} \right) + \dots \right] , \quad (14)$$

The residue of the propagator at the pole is

$$Z = 1 - \frac{1}{F^2 a^2} \left(I - \frac{1}{8} \right) + \dots . \quad (15)$$

Let us do the same with an arbitrary parameterization. Define new fields ϕ^a so that

$$\pi^a = \phi^a + C\phi^a \frac{\phi^2}{F^2} + \dots . \quad (16)$$

Note that the choice $C = -1/6$ corresponds to the exponential parametrization $U = \exp\{i\tau^a \phi^a/F\}$.

When substituting (16) in (4), the quadratic terms are not changed and the terms $\sim \phi^4$ have the form:

$$S^{(4)} = \frac{a^2}{F^2} \sum_{n,\mu} \left[\left(2C + \frac{1}{4} \right) \phi_n^4 - \frac{1}{4} \phi_{n+e_\mu}^2 \phi_n^2 - 2C \phi_n^2 \phi_n^a \phi_{n+e_\mu}^a \right] + \frac{M_0^2 a^4}{8F^2} \sum_n (1 + 8C) \phi_n^4 . \quad (17)$$




Figure 2: Renormalization of F_π .

In a general parameterization, the contribution from the loop graph in the pion polarizational operator is

$$\begin{aligned} \Pi^{\text{Fig.1}}(k) = \frac{2}{a^2 F^2} \left\{ \sum_{\mu} [(1 + 20C)D(0) - \cos(k_{\mu}a)D(ae_{\mu}) - 10CD(ae_{\mu}) - \right. \\ \left. 10C \cos(k_{\mu}a)D(0)] + \frac{5}{4}(1 + 8C)M_0^2 a^2 D(0) \right\} = \\ \frac{1}{a^4 F^2}(1 + 10C) + \frac{1}{a^2 F^2} \left\{ IM_0^2 \left(\frac{3}{2} + 10C \right) + k^2 \left[I(1 + 10C) - \frac{1}{8} \right] \right\} + o\left(\frac{1}{a^2}\right). \quad (18) \end{aligned}$$

The contribution from the measure is

$$\Delta M_0^2[\text{measure}] = -\frac{1}{a^4 F^2}(1 + 10C), \quad (19)$$

i.e quartic divergences coming from the measure and additive divergent contribution from the graph in Fig.1 cancel each other in any parameterization.¹

Solving again the dispersion equation with the polarization operator (18), one can readily see that the pole position of the propagator does not depend on the parameterization parameter C to the order $\sim 1/F^2$ and is given by the relation (14). Nothing prevents, however, for the residue of the pion propagator to depend on the parameterization, and it does:

$$Z = 1 - \frac{1}{F^2 a^2} \left[(1 + 10C)I - \frac{1}{8} \right] + \dots. \quad (20)$$

Let us calculate now the renormalization of the effective chiral coupling constant F . The latter is defined as the residue of the correlator of flavor non-singlet axial currents $A_{\mu}^a(x)$ at the pion pole

$$\int d^4x e^{ikx} \langle A_{\mu}^a(x) A_{\nu}^b(x) \rangle \sim F^2 \frac{k_{\mu} k_{\nu}}{k^2 - M_{\pi}^2} \delta^{ab}. \quad (21)$$

Technically, it is somewhat more convenient to pick up the coefficient of the structure $\sim g_{\mu\nu}$ in the correlator; due to the chiral current conservation in the chiral limit, the coefficients of these two tensor structures are the same.

There is only one relevant 1-loop diagram depicted in Fig.2.

¹A particular choice $C = -1/10$ where no quartic divergences in mass appear whatsoever is especially convenient.

It involves the vertex $\langle A_\mu A_\nu | \pi\pi \rangle$ which can be found by "covariantizing" the derivatives in Eq.(1) by the rule $\partial_\mu U \rightarrow \partial_\mu U - i(A_\mu U + U A_\mu)$ with $A_\mu \equiv A_\mu^a t^a$.² We have $\Delta F^2 = -2D(0)$ and hence, in the chiral limit,

$$F_{phys}^2 = F_0^2 \left[1 - \frac{2I}{F_0^2 a^2} + \dots \right] \quad (22)$$

in any parameterization.

3 Two-loop calculations.

We will calculate here only the additive two-loop corrections $\sim 1/F^4 a^6$ in the pion mass in the chiral limit ($M_0^2 \equiv 0$) and show that, after taking into account carefully all the contributions (including the relevant contributions from the measure), the final result is zero. We will work in the Weinberg parameterization.

To do the calculation, we are in a position to expand the action (2) up to the terms π^6/F^4 which have the form:

$$S^{(6)} = \frac{a^2}{8F^4} \sum_{n,\mu} (\pi_n^6 - \pi_n^4 \pi_{n+e_\mu}^2) . \quad (23)$$

The relevant two loop graphs are depicted in Fig.3 where crosses stand for the measure factor. Extracting the 6-pion vertex from Eq.(23) and substituting the free lattice propagator (5), the contribution of the graph in Fig.3a in the pion polarization operator is:

$$\Pi^a(k^2) = \frac{5}{a^2 F^4} \sum_\mu [3D^2(0) - D^2(ae_\mu) - 2\cos(k_\mu a)D(0)D(ae_\mu)] . \quad (24)$$

Setting here $k = 0$ and substituting the propagators (7, 8) in the form (9), we find the additive divergent contribution in the mass:

$$\Delta(M_\pi^2)^{Fig.3a} = \frac{10}{a^6 F^4} \left(I - \frac{1}{32} \right) . \quad (25)$$

The graph in Fig.3b can be calculated using the 4-pion vertices from Eq.(4). Putting $k = 0$ from the very beginning, we obtain

$$\begin{aligned} & \Pi^{Fig.3b}(0) = \\ & -\frac{1}{4a^6 F^4} \sum_{\mu,\nu} \int_{-\pi}^{\pi} \frac{d^4 x d^4 y}{(2\pi)^8} \frac{5 - 10\cos(x+y)_\mu + 2\cos x_\mu \cos y_\nu + 3\cos(x+y)_\mu \cos(x+y)_\nu}{\sum_{\alpha,\beta,\gamma} (1 - \cos x_\alpha)(1 - \cos y_\beta)[1 - \cos(x+y)_\gamma]} \end{aligned} \quad (26)$$

²This is all very much parallel to conventional calculations in chiral theory with dimensional regularization [1] or when calculating the effects due to finite temperature [4] (see recent [5] for review) or to an external magnetic field [6].




Figure 3: Polarization operator in two loops.

which can be rewritten as

$$\Pi^{\text{Fig.3b}}(0) = -\frac{1}{a^6 F^4} \left(4I - \frac{3}{16} \right) - \frac{1}{2a^6 F^4} J \quad (27)$$

where

$$J = \int_{-\pi/a}^{\pi/a} \frac{d^4 x d^4 y}{(2\pi)^8} \left[\frac{1}{\sum_{\mu} (1 - \cos x_{\mu})} - \frac{1}{\sum_{\mu} (1 - \cos(x + y)_{\mu})} \right] \left[1 - \frac{2}{\sum_{\nu} (1 - \cos y_{\nu})} \right]. \quad (28)$$

Integrating over $d^4 x$ first and using the fact that the integration region covers the whole period of the cosine functions (so that the shift $x_{\mu} \rightarrow x_{\mu} + y_{\mu}$ does not change the value of the integral), we see that $J = 0$. The corresponding additive mass correction is given by the first term in (27):

$$(\Delta M_{\pi}^2)^{\text{Fig.3b}} = -\frac{1}{a^6 F^4} \left(4I - \frac{3}{16} \right). \quad (29)$$

The graphs in Fig.3c,d have the same structure as that of the 1-loop graph in Fig.1, only we have to substitute there the 1-loop correction to the scalar propagator rather than the tree expression. We obtain:

$$\Pi^{\text{Fig.3c,d}}(k^2) = \frac{2}{a^2 F^2} \sum_{\mu} [\Delta D(0) - \Delta D(ae_{\mu}) \cos k_{\mu} a] \quad (30)$$

where

$$\Delta D(x) = - \int \frac{d^4 q}{(2\pi^4)} \frac{e^{iqx} \Pi^{(1)}(q)}{[\frac{2}{a^2} \sum_{\mu} (1 - \cos q_{\mu} a)]^2} , \quad (31)$$

with $\Pi^{(1)}(q)$ being given by the sum of the contributions (6) and (12) at $M_0^2 = 0$:

$$\Pi^{(1)}(q) = \frac{2D(ae_{\mu})}{a^2 F^2} \sum_{\mu} [1 - \cos(q_{\mu} a)] = \frac{2(I - 1/8)}{a^4 F^2} \sum_{\mu} [1 - \cos(q_{\mu} a)] .$$

We obtain

$$\begin{aligned} \Delta D(0) &= -\frac{1}{a^4 F^2} I \left(I - \frac{1}{8} \right) , \\ \Delta D(ae_{\mu}) &= -\frac{1}{a^4 F^2} \left(I - \frac{1}{8} \right)^2 . \end{aligned} \quad (32)$$

Therefore

$$(\Delta M_{\pi}^2)^{\text{Fig.3c,d}} = \Pi^{\text{Fig.3c,d}}(0) = -\frac{1}{a^6 F^4} \left(I - \frac{1}{8} \right) . \quad (33)$$

Finally, the Fig. 3e corresponds to the expansion of the factor $1/U^0$ in the group measure (11) up to the terms $\sim \pi^4$ which effectively produce the following correction to action:

$$S^{(6)}[\text{measure}] = -\frac{1}{4F^4} \sum_n \pi_n^4 , \quad (34)$$

and, correspondingly, to the 4-pion vertex. The calculation gives

$$(\Delta M_{\pi}^2)^{\text{Fig.3e}} = \Pi^{\text{Fig.3e}}(0) = -\frac{5I}{a^6 F^4} . \quad (35)$$

Adding up the additive ultraviolet divergent contributions (25), (29), (33) and (35) together, we find that total sum is zero, so that a massless pion stays massless, indeed.

Let us comment on the other possible contributions in the pion polarization operator to the order $\sim 1/F^4$. They come from the 1-loop graph in Fig.1 where the 4-pion vertex is extracted not from the leading order lagrangian (1) but from the higher derivatives terms:

$$L^{(4)} = \frac{1}{4} l_1 [\text{Tr}\{\partial_{\mu} U^+ \partial_{\mu} U\}]^2 + \frac{1}{4} l_2 [\text{Tr}\{\partial_{\mu} U^+ \partial_{\nu} U\}]^2 + \dots \quad (36)$$

where $l_{1,2}$ are some phenomenological parameters (or, better to say, some infinite constants in the free chiral lagrangian which will be transformed into finite phenomenological parameters after renormalization). However, the 4-pion vertices following from Eq.(36) are quartic in momenta. One can be convinced that the vertex is zero for zero value of the external momentum and the corresponding contribution is proportional to the square of external momentum and affects only Z-factor, not the additive term in the mass renormalization.

4 Conclusions

The method of calculations with the lattice regularization developed here presents an alternative to the standard dimensional regularization method. It is very explicit and physical and, though the calculations on the 4-dimensional lattice are slightly more involved than the calculations in the continuous space of the dimension $4 - \epsilon$ (or, rather just less habitual), these complications are not so serious. The main distinction compared to the dimensional regularization method is that power ultraviolet divergences are not "swept under the carpet", but are seen explicitly. Thereby, the main realm of application of our method could be the problems where the structure of power ultraviolet divergences in a non-renormalizable theory needs to be analyzed.

We calculated explicitly quadratically divergent pieces in F_π , M_π and in the residue of the propagator Z [see Eqs. (22), (14) and (15)]. We did not study the logarithmically divergent pieces whose structure should be roughly the same as with dimensional regularization. It would be an interesting methodic problem to verify it explicitly.

We are indebted to H. Leutwyler for illuminating discussions. This work was done under the partial support of RFBR INTAS grants 93-0283, 94-2851, RFFI grant 97-02-16131, CRDF grant RP2-132, and Schweizerischer National Fonds grant 7SUPJ048716.

References

- [1] J. Gasser and H. Leutwyler, *Annals Phys.* **158** (1984) 142; *Nucl. Phys.* **B250** (1985) 465.
- [2] P. Hasenfratz, *Phys. Lett* **141B** (1984) 385; W. Bietenholz, *Helv.Phys.Acta* **66** (1993) 633.
- [3] M. Creutz, *Quarks, gluons, and lattices* (Cambridge University Press, New York, 1983).
- [4] J. Gasser and H. Leutwyler, *Phys. Lett.* **184B** (1987) 837.
- [5] A.V. Smilga, *Phys.Repts.* **291** (1997) 1.
- [6] I.A. Shushpanov and A.V. Smilga, *Phys. Lett.* **402B** (1997) 351.